

## Conditional probability

How can we calculate the result in a case where two events are not independent. It means that, if one event occurs it will directly affect the probability for the other event?

If event A and B are those kind of complex events which will not exclude each other. In this case we have a so-called conditional probability (event A affects event B).

Notation:  $(p(A | B))$

In this case we mean the relative frequency which compares the sum of all probability to the probability of event B (probability of its occurrence).

$$p(A|B) = \frac{k_{AB}}{k_B} = \frac{\frac{k_{AB}}{k}}{\frac{k_B}{k}} = \frac{p(A \cap B)}{p(B)}$$

So we can get to the conclusion:

$$p(A \cap B) = p(A|B) p(B)$$

1.)  $(p(A \cap B))$ : This represents the probability that both events A and B occur simultaneously. It is also known as the probability of the intersection of A and B.

2.)  $(p(A|B))$ : This is the conditional probability of event A occurring given that event B has already occurred. It tells us how likely A is to happen under the condition that B has happened.

### What the Formula Says?

The formula states that the probability of both events A and B occurring together, is equal to the probability of B occurring multiplied by the probability of A occurring given that B has already occurred.

From:

<https://edu.iit.uni-miskolc.hu/> - Institute of Information Science - University of Miskolc

Permanent link:

[https://edu.iit.uni-miskolc.hu/tanszek:oktatas:techcomm:conditional\\_probability?rev=1724695271](https://edu.iit.uni-miskolc.hu/tanszek:oktatas:techcomm:conditional_probability?rev=1724695271)

Last update: 2024/08/26 18:01

